

CONTENTS

SL. No.	Title	Page No.
1	Title page	i
2	Dedication	ii
3	Certificate	iii
4	Declaration	iv
5	Acknowledgement	v - vi
6	List of symbols and abbreviations	vii-xiv
7	List of tables	xv-xvi
8	List of figures	xvii-xxiii
9	Contents	xxiv -xxxvi
10	Abstract	xxxvii- xxxviii

Chapter 1: Introduction	Page no
1.1. General Introduction	1-13
1.1.1. Medicinal plant and herbal utilization	3
1.1.2. Antioxidant agents from medicinal plants	4
1.1.3. Anti-inflammatory agents from medicinal plants	6
1.1.4. Anticancer agents from medicinal plants	7
1.1.5. Apoptosis	9
1.2. Plant Profile of <i>Calotropis gigantea</i> Linn. and literature review	13
1.2.1. Geographical distribution and description of the plant	
1.2.2. Taxonomy	15
1.2.3. Vernacular Names	15
1.2.4. Traditional use of <i>Calotropis gigantea</i>	15
1.2.5. Medicinal properties of <i>Calotropis gigantea</i>	16
1.2.6. Phytochemistry	
Chapter 2: Collection, Identification, Extraction, Chemical Screening and Characterization of <i>Calotropis gigantea</i> Linn.	19-36
2.1. Introduction	39 - 40
2.2. Materials and methods	41- 47
2.2.1. Chemicals	
2.2.2. Collection , identification and extraction of plant material	
2.2.3. Phytochemical screening of the extracts	
2.2.3.1. Tests for flavonoids	

- 2.2.3.2. Tests for alkaloids
- 2.2.3.3. Tests for steroids (Liebermann-Burchared test)
- 2.2.3.4. Test for triterpenoids (Salkowski test)
- 2.2.3.5. Test for reducing sugar
- 2.2.3.6. Test for gum (Molish's Test)
- 2.2.3.7. Test for tannins
- 2.2.3.8. Test for saponins
- 2.2.3.9. Tests for glycoside
- 2.2.3.10. Tests for cardiac glycosides (Keller-Killani test)
- 2.2.3.11. Test for resins
- 2.2.3.12. Tests for amino acid

2.2.4. Characterization by Analytical methods

- 2.2.4.1. IR spectroscopy
- 2.2.4.2. LCMS
- 2.2.4.3. NMR spectroscopy

2.3. Results

2.3.1. Phytochemical analysis

2.3.2. LCMS analysis of *C. giganteal* atex extract

- | | |
|--|-------|
| 2.3.2.1. LCMS analysis of EECGL | 48-67 |
| 2.3.2.2. LCMS analysis of WECGL | |
| 2.3.2.3. NMR analysis of <i>C. gigantea</i> latex extracts | |

2.4. Discussion

2.5. Conclusion

2.6. References

67-70

70

72-74

**Chapter 3:Chapter 3- Toxicity study of *Calotropis gigantea* Linn
latex extracts**

3.1.	Introduction	
3.2.	Mater	77-81
		81-94
ials and methods		
3.2.1.	Chemicals	
3.2.2.	Collection and extraction of plant material	
3.2.3.	<i>In-vitro</i> acute toxicity study	
3.2.3.1.	Brine shrimp lethality test Collection and hatching of brine shrimp	
3.2.3.2.	Collection and hatching of brine shrimp	
3.2.3.3.	Procedure of brine shrimp lethality bioassay	
3.2.3.4.	<i>In-vitro</i> toxicity study on the morphology of zebrafish embryos	
3.2.3.5.	Collection and hatching of zebra fish embryo	
3.2.3.6.	Procedure of toxicity study on zebra fish embryo	
3.2.4.	Toxicity study on human lymphocytes	
3.2.4.1.	Collection and separation of human lymphocytes	
3.2.4.2.	Culture of human lymphocytes	
3.2.4.3.	Estimation of reduced glutathione (GSH) in human lymphocytes	
3.2.4.4.	Estimation of Intracellular ROS measurement	
3.2.4.5.	DNA fragmentation study by alkaline comet assay	
3.2.5.	Sub-acute toxicity study in mice	
3.2.5.1.	Animals	
3.2.5.2.	Treatment schedule	
3.2.5.3.	Organ weight	
3.2.5.4.	Blood sample collection	

- 3.2.5.5. Red blood cell (RBC) count
- 3.2.5.6. White blood cell (WBC) count
- 3.2.5.7. Determination of haemoglobin
- 3.2.5.8. Estimation of urea
- 3.2.5.9. Estimation of creatinine
- 3.2.5.10. Measurement of serum alkaline phosphatase (ALP)
- 3.2.5.11. Measurement of serum glutamate oxaloacetate transaminase (SGOT) and serum glutamate pyruvate transaminase (SGPT)
- 3.2.5.12. Estimation of blood glucose
- 3.2.5.13. Measurement of serum cholesterol
- 3.2.5.14. Measurement of serum total protein
- 3.2.5.15. Histopathological study
- 3.2.6. Statistical analysis

3.3. Results

- 3.3.1. *In-vitro* acute toxicity study
 - 3.3.1.1. Effect of EECGL and WECGL on Brine shrimp lethality bioassay 94-108
 - 3.3.1.2. Effect of EECGL and WECGL on morphological toxicity in zebra fish embryos
 - 3.3.1.3. Effect of EECGL and WECGL on human lymphocytes
 - 3.3.1.3.1. Redox status (GSH levels)
 - 3.3.1.3.2. Intracellular ROS level
 - 3.3.1.3.3. Genotoxicity of EECGL and WECGL on human lymphocyte cells
 - 3.3.2. *In-vivo* sub-acute toxicity study
 - 3.3.2.1. Effect of ECGL and WECGL on organ weights
 - 3.3.2.2. Haematological parameters analysis

3.3.2.3. Effect of EECGL and WECGL on hepatic and renal biomarkers of mice	
3.4. Discussion	
3.5. Conclusion	
3.6. References	
 Chapter 4:Evaluation of <i>in-vitro</i> antioxidant and anti-inflammatory activity of Ethanol (EECGL) and Water extract (WECGL) of <i>Calotropis gigantea Linn</i> latex	
	108-113
	113
	115-122
1.Introduction	
4.1.1. Natural anti-oxidants	
4.1.2. Inflammation and anti-inflammatory agents	
4.2. Materials and methods	
4.2.1 Chemicals	
4.2.2 Collection of plant material and preparation of extracts	
4.2.3 <i>In-vitro</i> antioxidant activity	
4.2.3.1 Nitric oxide scavenging activity	125-130
4.2.3.2 DPPH scavenging activity	
4.2.3.3 Hydroxyl radical scavenging activity	131-140
4.2.3.4 Hypochlorous acid scavenging activity	
4.2.3.5 Superoxide anion scavenging activity	
4.2.3.6 Lipid peroxidation scavenging activity	
4.2.3.7 Peroxynitrite scavenging activity	
4.2.3.8 Determination of total phenolic content	
4.2.4 Anti-inflammatory activity	
4.2.4.1. <i>In-vitro</i> anti-inflammatory activity	
4.2.4.1.1. Human red blood cell (HRBC) membrane	
stabilization assay	
4.2.4.1.2. Inhibition of Protein denaturation	
4.2.4.2. <i>In-vivo</i> anti-inflammatory activity	
4.2.4.2.1. Animals	
4.2.4.2.2. Treatment Schedule	
4.2.4.2.3. Carrageenan-induced edema in rats	
4.2.5. Statistical analysis	
4.3. Results	
4.3.1. <i>In-vitro</i> antioxidant activities	

4.3.2. *In-vitro* anti inflammatory activities

4.4. Discussion

4.5. Conclusion

4.6. References

**Chapter 5- *In-vitro* cytotoxicity and apoptosis induction by
Calotropis gigantea Linn. latex in Jurkat cells and antimitotic
potential in *Allium cepa* root tip cells.**

5.1 Introduction

5.1.1 Leukaemia

5.1.2 Mitosis

5.1.3 Antimitotic drugs from phytochemicals and their
antimitotic potential on *Allium cepa*

140-149

5.2 Materials and methods

5.2.1 Chemicals

149-152

5.2.2 Collection of plant material and preparation of extracts

152-153

5.2.3 Cell lines

154-160

5.2.4 Isolation of human lymphocyte cell (HLC)

5.2.5 Experimental design of *in vitro* anticancer study on Jurkat
cells

5.2.6 *In-vitro* cytotoxic study

5.2.7 Measurement of intracellular reactive oxygen species
(ROS)

5.2.8 Chromatin condensation by propidium iodide and DAPI
staining

163-167

5.2.9 DNA fragmentation study by agarosegel electrophoresis

5.2.10 Antimitotic activity

5.2.10.1 Study of mitotic index

5.3 Results

- 5.3.1 EECGL and WECGL showed *in vitro* cytotoxicity on Jurkat cells 167-171
- 5.3.2 Intracellular ROS level
- 5.3.3 EECGL and WECGL produced chromatin condensation in Jurkat cell
- 5.3.4 EECGL and WECGL- induced apoptosis as measured by DNA laddering
- 5.3.5 Antimitotic effect of EECGL and WECGL

5.4 Discussion

5.5 Conclusion

5.6. References

Chapter 6- *In vitro and in vivo anticancer activity of ethanol (EECGL) and water extract (WECGL) of Calotropis gigantea Linn. Latex against Dalton ascites lymphoma (DLA) cells*

171-177

6.1 Introduction

6.2 Materials and methods

- 6.2.1 Chemicals and reagents
- 6.2.2 Animals
- 6.2.3 Cell lines
- 6.2.4 Isolation of mice lymphocyte cell (MLC)
- 6.2.5 *In-vitro* study
- 6.2.5.1** Experimental design for in-vitro study
- 6.2.5.2** *In-vitro* cell viability assay
- 6.2.5.3** Estimation of reduced glutathione (GSH)
- 6.2.5.4** Intracellular Reactive oxygen species (ROS)
 measurement
- 6.2.5.5** Nitric oxide release and generation assay
- 6.2.5.6** Lactate dehydrogenase release assay
- 6.2.5.7** Cellular morphology study by polarizing
 microscopy
- 6.2.5.8** Chromatin condensation by PI and DAPI
 staining
- 6.2.5.9** Acridine orange–ethidium bromide staining

177-179

6.2.5.10	DNA fragmentation study by agarosegel electrophoresis	179
6.2.5.11	Measurement of Mitochondrial membrane potential by Rhodamine 123	181-185
6.2.5.12	Study of DNA fragmentation by alkaline comet assay	
6.2.5.13	Detection of cellular apoptosis by flow cytometry	
6.2.6	<i>In-vivo</i> study	
6.2.6.1	Treatment schedule	188-191
6.2.6.2	Change in Body weight	191-203

6.2.6.3.	Studies on host survival time	
6.2.6.4.	Tumor volume	
6.2.6.5.	Tumor cell count	
6.2.6.6.	Determination of haemoglobin	
6.2.6.7.	Estimation of malondialdehyde (MDA) content	
6.2.6.8.	Estimation of reduced glutathione (GSH)	
6.2.6.9.	Estimation of superoxide dismutase (SOD)	
6.2.6.10.	Estimation of catalase (CAT)	
6.2.6.11.	Estimation of glutathione peroxidase (GP _X)	
6.2.6.12.	Estimation of glutathione-s-transferase (GST)	
6.2.7.	Statistical analysis	
6.3.	Results	204-226
6.3.1.	EECGL and WECGL suppress the <i>in vitro</i> cell proliferation of DLA cells	
6.3.2.	Cellular redox status (GSH levels).	
6.3.3.	Intracellular ROS level	
6.3.4.	Nitric oxide (NO) release and generation level	
6.3.5.	Lactate dehydrogenase (LDH) release level	
6.3.6.	EECGL and WECGL induce apoptosis on DLA cells	
6.3.7.	Effect of EECGL and WECGL on tumor Growth of DLA-bearing mice	
6.3.8.	Effect of EECGL and WECGL on haematological parameters	
6.3.8.1.	Effect of EECGL and WECGL on antioxidant parameters	
6.4.	Discussion	226-231
6.5.	Conclusion	
6.6.	References	231
		233-241

Chapter 7- *Calotropis gigantea* latex extracts exhibit antineoplastic potential against Ehrlich Ascites Carcinoma (EAC) cells through the induction of apoptosis and oxidative stress

7.1 Introduction	244-253
7.1.1 Etiology of Cancer	
7.1.2 Apoptosis	
7.1.3 Ehrlich ascites carcinoma (EAC) cell	
7.1.4 Alternative approaches in cancer therapy	
7.2 Materials and methods	
7.2.1 Chemicals and reagents	
7.2.2 Animals maintenance	253-261
7.2.3 Cell lines	
7.2.4 Mice lymphocyte cell (MLC)	
7.2.5 Experimental design for <i>in vitro</i> study	
7.2.6 <i>In vitro</i> study	
7.2.6.1 <i>In vitro</i> cell viability	
7.2.6.2 Oxidative stress parameters	
7.2.6.3 Nitric Oxide release and generation	
7.2.6.4 Lactate dehydrogenase release assay	
7.2.6.5 Lactate dehydrogenase release assay	
7.2.6.6 Cellular morphology study by scanning electron microscopy (SEM)	
7.2.6.7 Study on apoptosis	
7.2.6.8 . Detection of apoptosis by Annexin V-FITC Apoptosis detection kit	
7.2.6.9. Western blot analysis	
7.2.7. <i>In-vivo</i> anticancer activities	
7.2.7.1. Treatment schedule	
7.2.7.2. Body weight changes	
7.2.7.3. Tumor volume	
7.2.7.4. Red (RBC) and White (WBC) blood cell count	
7.2.7.5. Determination of Haemoglobin percentage	
7.2.7.6.	Det
ermination of malondialdehyde (MDA) content, reduced glutathione (GSH), and activities of superoxide dismutase (SOD), catalase, glutathione peroxidase (GPx) and glutathione s-transferase (GST) in EAC bearing mouse	
7.2.7.7. Histological study	
7.2.7.8. Study on peritoneal angiogenesis	

7.2.7.9.	Immunohistochemistry (IHC) study	
7.2.8.	Statistical analysis	
7.3.	Results	
7.3.1.	EECGL and WECGL induce the <i>in vitro</i> cytotoxicity in EAC cells	
7.3.2.	Cellular redox status (GSH levels).	261-284
7.3.3.	Intracellular ROS level	
7.3.4.	Nitric oxide release and generation level	
7.3.5.	Lactate dehydrogenase (LDH) release level	
7.3.6.	EECGL and WECGL drive apoptosis in EAC cells EECGL and WECGL treatment increased Bax and caspase-3	
7.2.7	EECGL and WECGL treatment increased Bax and caspase-3 expression but decreased Bcl-2 expression	
7.2.8	Effect of EECGL and WECGL on tumor regression of EAC bearing mice	
7.2.9	Effect on haematological parameters	
7.2.10	Effect on antioxidant parameters	
7.2.11	Anti angiogenic and antineoplastic effects of EECGL and WECGL	
7.3	Discussion	
7.4	Conclusion	285-291
7.5	References	291-292
Chapter 8: Summary and Conclusion		293-303
		304-310

Publications

Published papers

